

FoodData Central

Foundation Foods

Documentation and User Guide

(VERSION 3.29.19)

MARCH 2019

U.S. Department of Agriculture
Agricultural Research Service
Beltsville Human Nutrition Research Center
10300 Baltimore Avenue
Building 005, Room 107, BARC-West
Beltsville, Maryland 20705

Suggested Citation

U.S. Department of Agriculture (USDA), Agricultural Research Service. FoodData Central: Foundation Foods. Version Current: March 2019. Internet: www.fdc.nal.usda.gov.

Disclaimers

Reference to any product, service, process, or method by trade name, trademark, service mark, manufacturer or otherwise is for ease of identification only and does not imply recommendation, endorsement, or approval by, or an association with, the U.S. Department of Agriculture (USDA).

USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (e.g., Braille, large print, audiotape) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Issued March 2019

Contents

Preface	4
1. Introduction	5
1.1 About Foundation Foods Found in FoodData Central	5
1.2 Foundation Foods Highlights	5
1.2.1 Updated Nutrient Profiles	5
1.2.2 Expanded Information on Foods	5
2. Details on Information in Foundation Foods	6
2.1 Food Descriptions	6
2.2 Nutrient Data	7
2.2.1 Proximates	7
2.2.2 Minerals	8
2.2.3 Vitamins	8
2.2.4 Lipid Components	10
2.3 Weights	12
3. Reports and Data Files	13
3.1 Foundation Food Search Results	13
3.2 Overview of Data Tables	13
References	14

Preface

The food supply, and the scientific understanding of relationships between dietary intakes and health, have evolved over the years. USDA's food composition data resources also have evolved to meet the needs of diverse users, including researchers, nutrition professionals, health care providers, product developers, and policy makers. In recent years, the rapidly escalating pace of change in the food supply and the growing variety of uses for food data have greatly enhanced the need for transparent and easily accessible information about the nutrients and other components of foods and food products. FoodData Central is USDA's response to this need.

This integrated data system contains—in one place—five distinct types of food and nutrient composition data, each with a unique purpose. Three of these data types are well-established and familiar to many users: the **National Nutrient Database for Standard Reference**, the **Food and Nutrient Database for Dietary Studies**, and the **USDA Global Branded Food Products Database**.

The other two data types—**Foundation Foods** and **Experimental Foods**—represent “a bridge to the future” in food and nutrient composition. **Foundation Foods** includes nutrient values as well as extensive underlying metadata on commercially available foods. Foundation Foods data can provide valuable insights into the many factors that influence variability in nutrient profiles, and the number of foods in this data type will grow over time. **Experimental Foods** currently links to relevant agricultural research data from multiple sources, such as the [Agricultural Collaborative Research Outcomes System \(AgCROS\)](#). In future versions of FoodData Central, this data type will include information from multiple sources about foods that have been produced under experimental conditions and are not commercially available. The agricultural data in Experimental Foods will allow users to examine a range of factors, such as geography and agricultural practices, that may affect the nutritional profiles of foods and resulting dietary intake.

1. Introduction

1.1 About Foundation Foods Found in FoodData Central

Foundation Foods is a new food composition data type in the U.S. Department of Agriculture's (USDA) FoodData Central system. Foundation Foods contains expanded food nutrient profiles and metadata on commercially available foods, including the individual data points behind the mean values and metadata that include the number of samples, the location and dates on which samples were obtained, analytical methods used, and if appropriate, agricultural information such as genotype and production practices. These data allow users to see the variability in the nutrient values provided as well as the potential effects of the production site and procedures, season and climate, post-harvest processing, analytical methods, and other factors. The enhanced clarity and transparency around the data on these foods represent the future of FoodData Central and the number of these foods will grow over time.

Foundation Foods concentrates on basic foods (e.g., fruits, vegetables, flours) and staple foods (e.g., bread) eaten whole or used as ingredients. Some of the data have been acquired through the historical National Food and Nutrient Analysis Program (NFNAP) foods program. Other data will be from market and/or agriculturally acquired foods.

1.2 Foundation Foods Highlights

1.2.1 Updated Nutrient Profiles

Macronutrients (protein, fat, and carbohydrate) are the energy sources in the human diet. The inaugural version of Foundation Foods includes foods that appear in the final release of the National Nutrient Database for Standard Reference (SR Legacy) as well as new foods based on new acquisitions and analyses. For example, 68 of the foods in this data type have come from SR Legacy; the remaining items, which are the dry beans, are new acquisitions. Nutrient profiles taken from SR Legacy have been reworked to communicate full acquisition and analysis information, and thus, may vary slightly from those in SR Legacy. Additionally, data collected before the year 2000 have not been included, as those samples may include foods that are no longer representative of the marketplace. Agricultural sources will include fruits, vegetables, legumes, cereal grains, meats and dairy products, honey, and farmed fish.

1.2.2 Expanded Information on Foods

A key feature of Foundation Foods is the ability to see the specific value associated with each independent sample and hence the variability of the analyzed values for each component. For samples obtained at retail locations, Foundation Foods contains metadata on sample acquisition including city and state of purchase or manufacture, purchase date, expiration date (if applicable), product lot number, and UPC code (for retail products). For samples obtained from agricultural locations, metadata include information such as location (GPS coordinates), genotype, weather, and agricultural practices (e.g., conventional or organic farming) and analytical methodology.

Previously, NFNAP samples were generally collected in 12 locations based on a probability-proportional-to-size food sampling plan. For special cases, the sampling locations were significantly expanded to account for expected higher variability in nutrients (e.g., the sampling locations for drinking water were significantly expanded to account for the expected higher variability in minerals and fluoride). To maximize limited resources as well as the number of foods analyzed, the 12 samples were frequently composited with two randomly selected locations in each composite, for a total of 6 composites. Aliquots from these composites were sent to USDA qualified laboratories and collaborators for analysis. Although this approach generated useable mean values, sample-to-sample variability was lost. Therefore, the generated statistical parameters reflected the variability of the analytical samples, not the individual samples composited for analysis. Historically, the goal of the analyses was to generate a reliable mean across up to 150 nutrients for highly consumed foods. Moving forward, newer data as well as future data will be reported for independent samples analyzed from a single acquisition. In some cases, multiple samples from the same location may be needed to supply sufficient material for analysis. These will be composited and treated as a single independent sample.

2. Details on Information in Foundation Foods

The data for Foundation Foods are organized into three major categories: Food Descriptions, Nutrient Data, and Weights. These reflect the earlier approach to providing nutrient profile data but may change as FoodData Central evolves. In addition, support files are included that contain supplemental information related to these categories. File formats and related information are contained in the [Download & API Field Descriptions](#), available on the FoodData Central website. Abbreviations used in describing Foundation Foods are listed in Appendix B.

Data for agricultural products may be presented in forms not typically consumed but still representative of foods in the U.S. food supply. For example, data for common dry beans of different cultivars, growing locations, and climate conditions are presented on a 0% moisture basis rather than as sold.

2.1 Food Descriptions

The Food Descriptions category provides a full description of each food, including the name of the food, the brand name (if applicable), as well as the food's characteristics (e.g., raw or cooked, enriched or not, and color). Other fields in the Food Description file include:

- Scientific name.
- Common name, including alternative names for the product (e.g., soda or pop for a carbonated beverage), Uniform Retail Meat Identity Standard identification numbers, and USDA commodity codes as appropriate.
- Identification of food groups (see the [Download & API Field Descriptions](#) for more details) based on assignments in SR Legacy. These groupings are currently maintained to provide historical reference and continuity. It is anticipated that a current investigation in the area of ontology will result in changes in the food grouping systems.

- Amounts and physical descriptions, where appropriate, of refuse (inedible materials, such as seeds, bone, and skin); refuse amounts are expressed as a percentage of the total weight of the item as acquired and are used to compute the weight of the edible portion; most of the refuse data are obtained from measurements made for NFNAP samples.
- Factor used to calculate protein content from nitrogen content.
- Factors used to calculate number of kilocalories (kcal) from protein, fat, and carbohydrate, by difference.

Footnotes are provided for a few items where information about food description could not be accommodated in existing fields.

For details on the format of the Food Description file, see [Download & API Field Descriptions](#).

2.2 Nutrient Data

All nutrient values in Foundation Foods are based on analyses conducted by USDA under NFNAP or provided by other USDA units or external organizations. A unique code or FDC_ID number identifies individual foods contained in each of the data types. Currently, an FDC_ID number is assigned randomly when new or updated versions of foods are published in FoodData Central. However, when market and agricultural acquisition foods are presented through a Foundation Food, their FDC_ID are labeled as FDC Source ID to better distinguish them as sources for the current Foundation Food. The structure of this information is described in [Download & API Field Descriptions](#). Details on each of these analyses can be accessed through the documentation available with each dataset's download or through the "drill-down" capabilities (i.e., the ability to move from general information about a food to more detailed information) on the FoodData Central web site.

The values presented are calculated from analytical values, i.e., energy, protein, and vitamin A (i.e., retinol activity equivalents [RAE]) or they are rounded for display after the calculations are made. The source values, including protein, fat, and carbohydrates for energy, nitrogen for protein, and individual carotenoids and retinol for vitamin A, are stored in the data files to allow calculation of other nutrients and may have more decimal places than are presented in the reports and data files. A similar situation occurs when kilojoules (kJ) are calculated from kcal. Therefore, very small differences may occur if the user calculates from the rounded display values.

2.2.1 Proximates

"Proximate component" refers to the following macronutrients: water (moisture), protein, total lipid (fat), total carbohydrate, and ash. Except for a few food items, nutrient profiles contain values for the proximate components and at least one other nutrient.

The values for protein are calculated from the amount of total nitrogen in the food using the nitrogen-to-protein conversion factors recommended by Jones (1941) for most food items. If a specific factor is not available, the default value of 6.25 is used for the nitrogen-to-protein conversion factor.

The factor applied to each food item is provided in the NFactor field in the Food Description table. This differs from the approach taken in SR Legacy, which denotes protein as “analytical” if a laboratory has provided the value. Therefore, the protein value will be described differently between SR Legacy and Foundation Foods.

Total lipid values used on food labels represent the amount of triglyceride that would produce the amount of lipid fatty acids determined using gas chromatography, as required by the Nutrition Labeling and Education Act of 1990 (NLEA). The term “NLEA fat” is commonly used to refer to total fatty acids expressed as triglycerides.

Carbohydrate content, referred to as “carbohydrate by difference” in the tables, is expressed as the difference between 100 and the sum of the percentages of water, protein, total lipid (fat), ash, and alcohol (when present). Values for carbohydrate by difference include total dietary fiber content. “Sugars, total NLEA” refers to the sum of the values for individual monosaccharides (galactose, glucose, and fructose) and disaccharides (sucrose, lactose, and maltose), which are those sugars analyzed for nutrition labelling. Because the analyses of total dietary fiber, total sugars, and starch content are conducted separately and reflect the analytical variability inherent in the measurement process, the sum of these carbohydrate fractions may not equal the carbohydrate-by-difference value or may even exceed it.

Food energy is expressed in kcal and kJ. One kcal equals 4.184 kJ. The data represent physiologically available energy, which is the value remaining after digestive and urinary losses are deducted from gross energy (Merrill and Watt, 1973). Most energy values are calculated using the default factors of 4, 9, and 4 for protein, fat, and carbohydrates, respectively. Calorie factors for protein, fat, and carbohydrates are included in the Food Descriptions table for many food items. For foods containing alcohol, a factor of 6.93 is used to calculate kcal/g of alcohol (Merrill and Watt, 1973).

2.2.2 Minerals

Individual values for mineral data are reported in the database and include boron, calcium, cobalt, copper, fluoride, iron, magnesium, manganese, molybdenum, nickel, phosphorus, potassium, selenium, sodium, sulfur, and zinc. In other cases, such as the values for fluoride, selenium, and other minerals in NFNAP, samples for drinking water, select beverages, and grain-based products, respectively, regional and national averages are presented and should be used as such, as concentrations in local foods and beverages may vary widely.

2.2.3 Vitamins

Vitamins reported in the database include ascorbic acid, thiamin, riboflavin, niacin, pantothenic acid, vitamin B₆, vitamin B₁₂, folate, choline, vitamin A, vitamin D, vitamin E, and vitamin K. Many of the values were obtained in small sample sizes, often of composited samples.

Folate

Foundation Foods currently provides folate values in micrograms (μg) of dietary folate equivalents (DFEs). Future FoodData Central updates may present different forms of folate, including folic acid, food folate, and total folate, for some foods. Foundation Foods' DFEs are provided in accordance with the Institute of Medicine (IOM's) *Dietary Reference Intakes (DRI) for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* report (IOM, 1998). DFE concentrations differ from total folate concentrations only for foods containing synthetic folic acid added for enrichment or fortification. The calculation of DFEs reflects the greater bioavailability of synthetic folic acid than of naturally occurring food folate. To calculate DFEs for any single food, separate values are needed for naturally occurring food folate and added synthetic folic acid. The calculation is as follows:

$$\mu\text{g DFE} = \mu\text{g food folate} + (1.7 \times \mu\text{g folic acid})$$

Choline

Total choline, free choline (Cho), glycerophosphocholine (GPC), phosphocholine (PCho), phosphatidylcholine (PtdCho), and sphingomyelin (SM) from releases 1 and 2 of the USDA Database for the Choline Content of Common Foods (USDA, 2008) as well as newer values determined since the publication of those tables have been incorporated into Foundation Foods. Because metabolic pathways exist for the interconversion of Cho, GPC, PCho, PtdCho, and SM, total choline content is defined as the sum of the contents of these choline-contributing metabolites. Betaine values are not included in the calculation of total choline because the conversion of choline to betaine is irreversible (Zeisel et al., 2003).

Vitamin A

Values for vitamin A in μg of RAEs and μg of retinol are reported. One μg RAE is equivalent to 1 μg all-*trans*-retinol, 12 μg all-*trans*- β -carotene, or 24 μg other provitamin A carotenoids. Vitamin A activity values in RAE are calculated from the content of retinol and individual carotenoids (β -carotene, α -carotene, and β -cryptoxanthin) using well-established factors (IOM, 2001). Content of individual carotenoids (β -carotene, α -carotene, β -cryptoxanthin, lycopene, and lutein plus zeaxanthin) is reported in Foundation Foods.

Vitamin D

Vitamin D values in Foundation Foods are provided in micrograms. The biological activity of vitamin D is 40 international units per microgram (IU/ μg). Where available, values for specific isomers of vitamin D are reported, but only in μg . Cholecalciferol (vitamin D₃) is the form that is naturally present in animal products and most commonly added to fortified foods. Ergocalciferol (vitamin D₂) is the form found in plants and is added to some fortified foods, such as soy milk. In Foundation Foods, vitamin D content is the sum of vitamin D₂ and vitamin D₃ concentrations. Vitamin D may also be present as 25-hydroxycholecalciferol in some foods, such as fish, meat, and poultry, and this value is reported when available. The biological activity of 25-hydroxycholecalciferol has not been definitively determined, so it is not included in calculations of total vitamin D activity.

Vitamin E

The *Dietary Reference Intakes (DRI) for Vitamin C, Vitamin E, Selenium, and Carotenoids* report (IOM, 2000) defines vitamin E as the naturally occurring form (*RRR*- α -tocopherol) and three synthetic forms of α -tocopherol. Foundation Foods provides vitamin E values in mg of α -tocopherol (nutrient 323) in accordance with this DRI report. Although β -, γ -, and δ -tocopherol do not contribute to vitamin E activity, they are included in Foundation Foods when analytical data are available.

Vitamin K

Data on vitamin K₁ (phylloquinone), dihydrophylloquinone, and menaquinone-4 are presented individually in Foundation Foods. Dihydrophylloquinone is a form of vitamin K that is created during the commercial hydrogenation of plant oils. Menaquinone-4 is formed from vitamin K₁ and/or the synthetic form of vitamin K in animal feed, and is found primarily in meats and meat products.

2.2.4 Lipid Components

Fatty acid values are expressed in g per 100 g of food. Often, there are more values for total lipid than for individual fatty acid values. Due to analytical variability, the sum of the fatty acids may not add up to the value for total lipid. Total lipid values used on food labels represent the amount of triglyceride that would produce the amount of lipid fatty acids determined using gas chromatography, as required by the NLEA.

The basic format for describing individual fatty acids is that the number before the colon indicates the number of carbon atoms in the fatty acid chain, and the number after the colon indicates the number of double bonds. For unsaturated fatty acids, additional nutrient numbers have been added to accommodate the reporting of many specific positional and geometric isomers. Of the specific isomers, two basic classifications are considered: omega double bond position and *cis/trans* configuration of double bonds.

Omega-3 (n-3) and omega-6 (n-6) isomers are denoted in shorthand nomenclature as n-3 and n-6. The n- number indicates the position of the first double bond from the methyl end of the carbon chain. The letter *c* indicates a *cis* bond, and the letter *t* indicates a *trans* bond. For polyunsaturated fatty acids, *cis* and *trans* configurations at successive double bonds may be indicated. For example, linoleic acid is an 18-carbon omega-6 fatty acid with two double bonds, both in *cis* configuration. When data are isomer specific, linoleic acid is described as 18:2 n-6 *c,c*. Other isomers of 18:2, for which nutrient numbers have now been assigned, include 18:2 *c,t*, 18:2 *t,c*, 18:2 *t,t*, 18:2 *t* not further defined, and 18:2 *i*. 18:2 *i* is not a single isomer but includes isomers other than 18:2 n-6 *c,c* with peaks that cannot be easily differentiated in the particular food item. Systematic and common names for fatty acids are provided in Table 1.

Table 1. Systematic and Common Names for Fatty Acids

Fatty acid	Systematic name	Common name of most frequent isomer
Saturated fatty acids		
4:0	Butanoic acid	Butyric acid
6:0	Hexanoic acid	Caproic acid
8:0	Octanoic acid	Caprylic acid
10:0	Decanoic acid	Capric acid
12:0	Dodecanoic acid	Lauric acid
13:0	Tridecanoic acid	
14:0	Tetradecanoic acid	Myristic acid
15:0	Pentadecanoic acid	
16:0	Hexadecanoic acid	Palmitic acid
17:0	Heptadecanoic acid	Margaric acid
18:0	Octadecanoic acid	Stearic acid
20:0	Eicosanoic acid	Arachidic acid
22:0	Docosanoic acid	Behenic acid
24:0	Tetracosanoic acid	Lignoceric acid
Monounsaturated fatty acids		
14:1	Tetradecenoic acid	Myristoleic acid
15:1	Pentadecenoic acid	
16:1 undifferentiated 16:1 <i>cis</i> *	Hexadecenoic acid	Palmitoleic acid
17:1	Heptadecenoic acid	
18:1 undifferentiated 18:1 <i>cis</i> *	Octadecenoic acid	Oleic acid
20:1	Eicosenoic acid	Gadoleic acid
22:1 undifferentiated 22:1 <i>cis</i> *	Docosenoic acid	Erucic/citoleic acid
24:1 <i>cis</i>	Cis-tetracosenoic acid	Nervonic acid
Polyunsaturated fatty acids		
18:2 undifferentiated 18:2 <i>i</i> (mixed isomers) 18:2 n-6 <i>cis, cis</i> * 18:2 conjugated linoleic acid	Octadecadienoic acid	Linoleic acid
18:3 undifferentiated 18:3 n-3 <i>cis, cis, cis</i> * 18:3 n-6 <i>cis, cis, cis</i> 18:3 <i>i</i> (mixed isomers)	Octadecatrienoic acid	Linolenic acid Alpha-linolenic acid Gamma-linolenic acid
18:4	Octadecatetraenoic acid	Parinaric acid
20:2 n-6 <i>cis, cis</i>	Eicosadienoic acid	
20:3 undifferentiated	Eicosatrienoic acid	

Fatty acid	Systematic name	Common name of most frequent isomer
20:3 n-3		
20:3 n-6		
20:4 undifferentiated		
20:4 n-6*	Eicosatetraenoic acid	Arachidonic acid
20:5 n-3	Eicosapentaenoic acid	Timnodonic acid
21:5		
22:4		
22:5 n-3	Docosapentaenoic acid	Clupanodonic acid
22:6 n-3	Docosahexaenoic acid	
Trans fatty acids		
Fatty acids, total trans-monoenoic		
16:1 <i>trans</i>		
18:1 <i>trans</i>		
22:1 <i>trans</i>		
Fatty acids, total trans-polyenoic		
18:2 <i>trans</i> not further defined		
18:2 <i>trans, trans</i>		

Isomer associated with the common name; the most frequent isomer is listed for the undifferentiated fatty acid. It is not possible to include every possible geometric and positional isomer in the database. Where specific isomers exist for a fatty acid, the common name of the most frequent isomer is listed for the undifferentiated fatty acid and an asterisk () designates the isomer to which that name applies. For example, the most frequent isomer for 18:1 is oleic acid. Therefore, undifferentiated 18:1 is designated in Table 1 as oleic acid, and the asterisk indicates that the common name for 18:1, oleic acid, only applies to this isomer.

Because cholesterol is found only in foods of animal origin, cholesterol values are provided only for foods of animal origin and foods containing at least one ingredient of animal origin (e.g., cake that contains eggs).

2.3 Weights

Portions are provided in grams for edible material without refuse (i.e., the edible portion of the food), such as an apple without the core or stem or a chicken leg without the bone. These determinations of edible portion are on a 100-gram or percent basis. Also provided is information on portion sizes for most food items (e.g., 1 cup, 1 tablespoon, 1 fruit, or 1 leg); all foods are reported in 100 g. The weights are determined from samples acquired as part of NFNAP. It should be noted that portions and weight are unique to each data type in FoodData Central—Foundation Foods, Food and Nutrient Database for Dietary Studies (FNDDS), USDA Global Branded Food Products Database, SR Legacy, and Experimental Foods. In some cases, there may be many more determinations of portion sizes than there are of the nutrient analyses. Although efforts have been made to provide representative values, portion sizes obtained from different sources vary considerably for some foods. Portions for specific foods are

displayed in the measure tab on the FoodData Central website or in the food portion file in the download files; the format of this file is described in the [Download & API Field Descriptions](#).

The gram weights in the food portion file can be used to calculate nutrient values for food portions from the values provided per 100 g of food. The following formula is used to calculate the nutrient content per portion:

$$N = (V*W)/100;$$

where:

N = nutrient value per portion size,

V = nutrient value per 100 g (Nutr_Val in the Nutrient Data file), and

W = weight (in g) of portion (Gm_Wgt in the Weight file).

3. Reports and Data Files

3.1 Foundation Food Search Results

Using the FoodData Central search program (www.fdc.nal.usda.gov), users can look up the nutrient content of any food in Foundation Foods as well as those in SR Legacy, USDA Global Branded Food Products Database, and FNDDS. Foods may be selected by key terms, such as nutrient name, food name, NDB# (in SR Legacy), FDC_ID number, or brand (in USDA Global Branded Foods Database).

An application program interface (API) is also available for developers to use to access the database with their own applications. They can be assured that they are linking to the most up-to-date version of the database. Details on using the API are provided on the FoodData Central web site (www.fdc.nal.usda.gov).

3.2 Overview of Data Tables

The data files for Foundation Foods are available from the FoodData Central web site (www.fdc.nal.usda.gov) in ASCII (ISO/IEC 8859-1), and Microsoft Access 2016 formats.

Descriptions of each field in these tables and the relationships between them are contained in the [Download & API Descriptions](#).

References

Institute of Medicine. Dietary Reference Intakes (DRI) for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline. Washington, DC: National Academy Press; 1998.

Institute of Medicine. Dietary Reference Intakes (DRI) for Vitamin C, Vitamin E, Selenium, and Carotenoids. Washington, DC: National Academy Press; 2000.

Institute of Medicine. Dietary Reference Intakes (DRI) for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Washington, DC: National Academy Press; 2001.

Merrill AL, Watt BK. Energy value of foods: basis and derivation, revised. Agriculture Handbook 74. Washington, DC: U.S. Department of Agriculture; 1973.

U.S. Department of Agriculture. USDA Database for the Choline Content of Common Foods. Release 2; 2008. Available at: www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/nutrient-data-laboratory/docs/usda-database-for-the-choline-content-of-common-foods-release-2-2008. Accessed January 4, 2019.

Zeisel SH, Mar MH, Howe JC, Holden JM. Concentrations of *choline*-containing compounds and betaine in common foods. *Journal of Nutrition*. 2003;133(5):1302-1307.